

Soluble ST2 as a Potential Biomarker in Pericardial Fluid of Coronary Artery Patients

Reşat Dikme¹, MD; Mahmut Padak¹, MD; Mesut Işık², MD; İsmail Koyuncu³, MD; Ebru Temiz⁴, MD; Mehmet Salih Aydın⁵, MD; Ömer Göç¹, MD

DOI: 10.21470/1678-9741-2020-0317

Abstract

Introduction: The growth Stimulation expressed gene 2 (ST2) (or interleukin 1 receptor-like 1, also known as IL1RL1) is considered a biomarker of poor prognosis in cardiovascular diseases. The aims of this study are to investigate ST2 in the pericardial fluid (PF) of coronary artery disease patients and to contribute to the understanding of the pathophysiology of coronary artery disease.

Methods: 40 patients (blood plasma and PF) who underwent coronary artery bypass surgery and 40 controls (blood plasma only) were included in this study. Soluble ST2 (sST2) level was determined by enzyme-linked immunosorbent assay method in plasma and PF, and sST2 gene expression was determined by quantitative real-time polymerase chain reaction (QRT-PCR) method.

Results: The sST2 level was found to be 44.89 ng/ml and 390.357

ng/ml in the control and patient groups' plasma, and 223.992 ng/ml in the PF of the patient group. An increase in sST2 level was detected in the patient group compared to the control group ($P<0.001$). The sST2 expression in plasma was higher in the patient group than in the control group. Additionally, sST2 was more expressed in the plasma of the patient group than PF ($P<0.001$).

Conclusion: The fact that sST2 was detected for the first time in a high level in PF showed that this biomarker was closely related with the heart and strengthened its potential to be used as a biomarker. Therefore, sST2 can contribute to the understanding of the pathophysiology of coronary artery disease.

Keywords: Coronary Artery Disease. Pericardial Fluid. Real-Time Polymerase Chain Reaction. IL1RL1 protein, human. Biomarkers. Prognosis.

Abbreviations, acronyms & symbols			
AVG	= Average	mRNA	= Messenger ribonucleic acid
CAD	= Coronary artery disease	PCR	= Polymerase chain reaction
cDNA	= Complementary deoxyribonucleic acid	PF	= Pericardial fluid
CPB	= Cardiopulmonary bypass	PHFS	= Penn Heart Failure Study
Ct	= Cycle threshold	RNA	= Ribonucleic acid
CVD	= Cardiovascular diseases	QRT-PCR	= Quantitative real-time polymerase chain reaction
DNA	= Deoxyribonucleic acid	sST2	= Soluble growth Stimulation expressed gene 2
ELISA	= Enzyme-linked immunosorbent assay	ST2	= Growth Stimulation expressed gene 2
FHS	= Framingham Heart Study	ST2L	= Membrane-bound growth Stimulation expressed gene 2 or mST2
GE	= Genomic elution	STEMI	= ST-elevation myocardial infarction
IL33	= Interleukin 33		

¹Department of Perfusion Techniques Program, Vocational School of Health Services, Harran University, Şanlıurfa, Turkey.

²Department of Bioengineering, Faculty of Engineering, Bilecik Seyh Edebali University, Bilecik, Turkey

³Department of Medical Biochemistry, Faculty of Medicine, Harran University, Şanlıurfa, Turkey.

⁴Department of Medical Biology and Genetics, Vocational School of Health Services, Harran University, Şanlıurfa, Turkey.

⁵Department of Cardiovascular Surgery, Faculty of Medicine, Harran University, Şanlıurfa, Turkey.

This study was carried out at the Harran University, Şanlıurfa, Turkey.

Correspondence Address:

Mesut Işık

 <https://orcid.org/0000-0002-4677-8104>

Department of Pharmacy Services, Bilecik Seyh Edebali University, Faculty of Engineering, Department of Bioengineering
Gülümbe, Bilecik, Turkey

Zip Code: 11230

E-mail: mesut.isik@bilecik.edu.tr

Article received on June 21st, 2020.

Article accepted on June 25th, 2020.

INTRODUCTION

The incidence of cardiovascular diseases (CVD), which is one of the most important causes of mortality and morbidity in the world, is rapidly increasing. Despite the ongoing developments in their treatment, these diseases are still in the first place as the cause of death in the People's Republic of China^[1].

A biomarker expressed in the heart tissue and coronary endothelial cells can be released into the blood and pericardial fluid (PF). Since some of the heart-related biomarkers detected in the circulation are expressed in both the heart muscle and skeletal muscle, analyses in the blood may not indicate heart disease. Therefore, besides blood, PF can be used for diagnosis in CVD. The PF within the pericardial cavity, which is the double-walled pouch enveloping the entire heart, shows dynamic changes. The pericardium has immunological, paracrine, vasomotor, and fibrinolytic activity and it affects the myocyte structure, function, and gene expression by synthesizing biomolecules such as eicosanoids (prostaglandin, thromboxane, and leukotriene)^[2]. Recently, the growth Stimulation expressed gene 2 (ST2) (or interleukin 1 receptor-like 1 [also known as IL1RL1], T1, Fit1, IL33R, DER4) has been considered as an important prognostic biomarker because of its involvement in the pathogenesis of CVD, its stability, and easy measurement in clinical samples. ST2, which plays an important role in cellular stress and inflammatory response, is the receptor of interleukin 33 (IL33)^[3]. IL33, with its

cytokine and transcriptional feature, creates a signal between cells as a nuclear factor and helps to protect the heart^[4,5]. IL33, which creates a paracrine effect between cardiac myocytes and fibroblasts when there is mechanical load in the heart, acts by binding to ST2^[4]. ST2 has two important isoforms, these isoforms are membrane-bound ST2 (ST2L or mST2) and soluble ST2 (sST2). ST2L is not free in peripheral circulation. sST2 without transmembrane and intracellular domain moves freely in the peripheral circulation^[5]. While binding of ST2L to IL33 has a cardioprotective effect (mechanical stimulation, decreased cardiac damage, prevention of apoptosis, lowered inflammatory effect, hypertrophy, and fibrosis), binding of sST2 to IL33 causes these positive effects to disappear (Figure 1). sST2 acts as a trap receptor (decoy receptor), reducing the positive effects of IL33^[4].

High level of sST2 indicates that the heart is under intense stress. This causes cellular death, tissue fibrosis, decreased heart function, and increased rate of disease progression. Therefore, sST2 is considered to be a biomarker of poor prognosis in CVD^[5]. It was found that sST2 rising above the normal level increased many negative conditions due to CVD by approximately three times^[6]. According to multiple clinical studies, sST2 has emerged as a clinically useful prognostic biomarker in patients with CVD, such as myocardial infarction, acute dyspnea, and heart failure, and in the low-risk population^[7].

The main source of sST2 in individuals with healthy and different diseases is currently not fully determined. In a study of




Fig. 1 – Cardioprotective relationship of interleukin 33 (IL33) with membrane-bound growth STimulation expressed gene 2 (ST2L) and blocking with soluble growth STimulation expressed gene 2 (sST2). DNA=deoxyribonucleic acid.

neonatal rat heart myocytes related to heart disease, the sST2 in the bloodstream was found to be of myocardial origin^[8]. However, some studies have shown that the predominant source of sST2 in cardiac patients may be vascular endothelial cells rather than the myocardium^[9]. When all the studies are evaluated together, it is revealed that sST2 is produced by cardiac fibroblast, cardiomyocytes, macrovascular (aorta and coronary arteries), and microvascular endothelial cells in case of damage and stress. In another study, parallel increased release of both ST2L and sST2 was detected in cardiomyocytes and cardiac fibroblasts under biomechanical stress^[10]. However, there are no studies on sST2 release in epicardium and pericardium. We assume that two of the sources of sST2 in humans are probably epicardium and pericardium. In coronary artery disease (CAD) patients, research of sST2 only in the blood is not sufficient. For this reason, it is also important to detect sST2 in the PF, the tissue closest to the heart. The fact is that the increase mechanism of ST2, which has a negative effect on the formation of CAD, is not known; there is no ST2 study in the PF in the literature, and unanswered questions about the IL33-ST2 pathway make this study important. Enzyme-linked immunosorbent assay (ELISA) kits are often used to measure human sST2 levels in serum and plasma. In this study, differently, gene expression study was performed for the measurement of sST2.

METHODS

Patients Included in the Study

The study protocol was carried out in accordance with the Helsinki Declaration as revised in 1989. Signed informed consent forms were obtained from the families of all participants. Ethical approval was obtained from the Harran University Local Research Ethics Committee (Ethical approval number: 05.01.2017-17/01/23). Forty patients who underwent coronary artery bypass surgery with cardiopulmonary bypass (CPB) method were included in the study. Both PF and blood were taken from the patient group. As for the control group, only blood samples were taken from 40 healthy individuals in the same age group.

Obtainment of PF

The pericardium was opened after median sternotomy was performed in patients undergoing coronary bypass surgery with the CPB method. Then, PF was aspirated with a sterile syringe. The aspirated PF was placed in sterile tubes without gel (10 ml, Vacutainers/BD Biosciences) and put in an ice-filled container. Then, the PF in the sterile tube was purified from the cells by passing through the centrifugation step at 5000 rpm for five minutes. After the centrifugation step, the supernatant was taken to a ribonuclease-free tube (1.5 mL, Eppendorf) and stored at -80 °C.

Obtainment of Blood Plasma

Blood was taken from the patients who underwent coronary bypass surgery with the CPB method, before CPB, and gel-free anticoagulant (0.2 ml, Nevparin injectable 25000 IU/5 mL) was placed in sterile glass tubes (10 ml, Vacutainers/BD Biosciences).

These tubes were immediately placed in an ice-filled container and delivered to the laboratory. Then, sterile tubes were centrifuged at 5000 rpm for five minutes. After the centrifugation step, plasma was taken into a ribonuclease-free tube (1.5 mL, Eppendorf) and stored at -80 °C. After collecting PF and blood from 40 patients, ELISA and quantitative real-time polymerase chain reaction (QRT-PCR) studies were performed.

Studying Stages of ST2 Gene Expression in PF and Plasma

1. Total ribonucleic acid (RNA) (messenger RNA [mRNA]) Isolation in PF and Plasma

For isolation of mRNA in PF and plasma, Qiagen (Qiagen GmbH, Hilden, Germany) miRNeasy Serum/Plasma Kit (Cat No./ID: 217184) was used. The working procedure of the manufacturer company was followed and total RNA isolation was performed.

2. Complementary deoxyribonucleic acid (cDNA) Obtainment from mRNA in PF and Plasma

In order to transfer mRNA in obtained total RNA to cDNA, RT2 HT First Strand Kit (Qiagen, GmbH, Hilden, Germany, Cat No./ID: 330411) was used. Each reverse transcribe phase was carried out twice. The contents of reverse transcriptase PCR reaction mixture applied to obtain cDNA from mRNA were prepared as in Table 1. According to the work flow chart of the kit, the relevant procedure was followed.

Table 1. Reverse transcriptase PCR mixture for obtaining cDNA from mRNA.

Ingredients	Amount
RNA	500 ng/μl
Buffer GE (removal of genomic DNA)	2 μl
Ribonuclease-free water	Adjusted to total 10μl

cDNA=complementary deoxyribonucleic acid;
DNA=deoxyribonucleic acid; mRNA=messenger ribonucleic acid; GE= Genomic elution; PCR=polymerase chain reaction; RNA=ribonucleic acid

3. Quality and quantity identification from cDNA obtained from PF and plasma

Quality and quantity identification was done by using Thermo Scientific Multiskan Spectrophotometer System (Waltham, Massachusetts, United States of America) equipment. Before expression analysis experiments, cDNA quantities obtained from samples were declined to 50 ng/μl for mRNA cDNA.

4. Real-time PCR (QRT-PCR) Method

Before QRT-PCR, mRNA-cDNA preamplification was performed in PF and plasma, and then routine PCR protocol was applied. Rotor Gene 6000 Real-Time PCR Machine (Qiagen GmbH, Hilden, Germany) device and QuantiTect SYBR Green PCR

Kit (Qiagen GmbH, Hilden, Germany, Cat No./ID: 204143) were used for QRT-PCR.

Primer pool: 40 μ l (20 μ l F [10 pmol/ μ l] + 20 μ l R [10 pmol/ μ l]) primer was used; 210 μ l of water was added per primer and the relevant content is given in Table 2.

The reaction was carried out at 25.5 μ l by adding 5 μ l cDNA to the total mix. Routine PCR procedure was applied according to Table 3 for sST2 expression level in PF and Plasma.

The product obtained after PCR was diluted with 91 μ l of water; 5 μ l was used for real time PCR (QRT-PCR). For QRT-PCR, the component and volume were used according to Table 4.

The prepared mixture was distributed to the tubes to be 9.5 μ l; 0.5 μ l cDNA of each sample was added to each tube containing the QRT-PCR premix and the study was performed under the PCR conditions set forth in Table 5.

5. Primer Design

In this study, ST2 primers (Metabion International AG/metabion GmbH, Germany) and beta-actin, a housekeeping gene, were used. ST2 primers were designed using the "Gene" interface (<http://www.ncbi.nlm.nih.gov/tools/primerblast/>) in the National Center for Biotechnology Information database. The sequences of the ST2 and beta-actin primers are shown in Table 6.

ST2 ELISA Study in PF and Plasma

Human sST2 ELISA Kit (Elabscience Biotechnology Co., Ltd., Product ID: E-EL-H1615) was used to determine the ST2 level in PF and plasma by ELISA method. Samples in PF and plasma were kept at room temperature for two hours before working. Then, sST2 levels were measured according to the manufacturer's operating procedure.

Statistical Analysis

After the studies were completed, it was checked whether the data were normally distributed. Then, one-way analysis of variance was performed. According to the Shapiro-Wilk test result, Student's *t*-test was used for the significant difference test in the normal distribution of the groups, and the Mann-Whitney U test was used in the abnormal distribution. In our study, Mann-Whitney U test was used for statistical analysis of sST2 ELISA results in PF and plasma, and Student *t*-test was used in sST2 gene expression results. By comparing each test group with the control groups, *P*-values <0.05 were considered significant. In our study, "prism 6" program (GraphPad Software, Inc, San Diego,

United States of America) and QIAGEN analysis program (<https://www.qiagen.com/us/shop/genes-and-pathways/data-analysis-center-overview-page/>) were used for statistical analysis.

Table 2. Primary pool mix for ST2 gene expression.

For 1 sample	
5' miScript master mix	12.5 μ l
Taq polimeraz	0.5 μ l (250 unit)
Primer pool	7.5 pmol/ μ l
Total mix: 20.5 μ l	

ST2=growth Stimulation expressed gene 2

Table 3. Routine PCR protocol for ST2 expression level.

Temperature ($^{\circ}$ C)	Time	
95	10 minutes	
94	15 seconds	} 12 times
60	2 minutes	
4		

PCR=polymerase chain reaction; ST2=growth STimulation expressed gene 2

Table 4. QRT-PCR components for sST2 expression level.

Components	Volume
RT ² SYBR [®] Green qPCR Mastermix	5 μ l
F Primer (10 pmol/ μ l)	0.3 μ l
R Primer (10 pmol/ μ l)	0.3 μ l
Ribonuclease/deoxyribonuclease-free water	3.9 μ l
cDNA (50 ng/ μ l)	0,5 μ l
Total	10 μ l

cDNA=complementary deoxyribonucleic acid; QRT-PCR=quantitative real-time polymerase chain reaction; sST2=soluble growth STimulation expressed gene 2

Table 5. QRT-PCR settings for ST2.

	Temperature ($^{\circ}$ C)	Time	
Enzyme activation	95	10 minutes	
Denaturation	95	15 seconds	} 40 times
Primer binding/elongation	60	60 seconds	

QRT-PCR=quantitative real-time polymerase chain reaction; ST2=growth STimulation expressed gene 2

Table 6. ST2 and beta-actin forward and reverse primers.

	Forward primer	Reverse primer
ST2	5'-TTCCAGTAATCGGAGCCCCT-3'	5'-AACCATGTGCAAGGGGAAGAG-3'
Beta-actin	5'-CGTACCACAGGCATTGTGATG-3'	5'-TTTGATGTCACGCACGATTTC-3'

ST2=growth STimulation expressed gene 2

RESULTS

As a result of our ELISA study, the sST2 level was found to be 223,992±19,628 ng/ml in PF, and 44,89±4,226 ng/ml and 390,357±25,245 ng/ml in plasma of the control and patient groups ($P<0.001$) (Figure 2). In our study, the fold changes of sST2 gene expression levels in PF and plasma were evaluated. The fold changes of sST2 gene expression levels in PF and plasma were evaluated with Student's *t*-test. As a result, the average Ct value for sST2 was found as 13.33 in PF, and 19.69 and 2.26 in the plasma of the patient and control groups, respectively. As a result, the average Delta (Ct)* value for sST2 was found as 0,790741 in PF, and 0,39037 and -0,7450 in the plasma of the patient and control groups, respectively. While $2^{22\Delta(-\text{Ort. Delta Ct})}$ value of sST2 was found as 0.578047 in PF, it was found as 1.31073 and 0.1506 in the patient and control groups' plasma, respectively. The sST2 fold change ($2^{-\Delta\Delta\text{Ct}}$) in the plasma of patients was calculated as 2.2675 compared to PF (Figure 3). As a result of statistical analysis, significant changes were found in the replicate $2^{\Delta(-\text{Delta Ct})}$ values of sST2 gene expression in PF and plasma ($P<0.001$).

DISCUSSION

In the American College of Cardiology Foundation/American Heart Association — ACCF/ AHA — guide, ST2 was defined as a “novel biomarker” in 2013^[11]. The sST2, a biomarker independent of natriuretic peptides and cardiac troponins, has been accepted as an important biomarker in many diseases such as ST-elevation myocardial infarction (STEMI), non-STEMI (or NSTEMI), CAD, heart failure, valvular diseases, cardiomyopathy, coronary bypass, acute cardiac allograft rejection, and acute Kawasaki disease^[12]. According to another study, sST2 shows itself as a strong ischemic marker and is used especially in predicting the etiologies of cardiomyopathy^[11]. Studies have reported that ST2 is not affected by factors such as age, body mass index, and renal dysfunction, and its *in vitro* stability is high^[13,14].

According to the Framingham Heart Study (FHS) Cohort analysis, the sST2 level gives important information in CAD asymptomatic individuals^[15].




Fig. 2 - Determination of soluble growth STimulation expressed gene 2 (sST2) level of as a biomarker for coronary artery disease (CAD) in pericardial fluid (PF) and plasma.




Fig. 3 - Fold change values of soluble growth STimulation expressed gene 2 in pericardial fluid (PF) and plasma of coronary artery disease.

The most important reference values for circulating sST2 were obtained from the FHS. While the reference range of ST2 level was 11-45 ng/mL in men (n=462), sST2 was found to be 9-35 ng/mL in women (n=674)^[15,6].

Dieplinger et al.^[6], while the normal mean value of sST2 was 18 ng/ml, stated that concentrations > 35 ng/ml were at high risk for CVD. In the Penn Heart Failure Study (PHFS), it was stated that high sST2 levels gave a predictive idea about death and heart transplantation in patients with heart failure^[16]. According to the PHFS, patients with sST2 value > 35 ng/ml were found to have 2.8 times more risk of CVD than normal individuals^[17].

Another study reported that heart patients with high sST2 values may have an average of 32% heart failure and 45% death after 11 years^[7]. The increase in the level of sST2 in patients with stable CAD may have been caused by chronic inflammation in atherosclerosis. According to large-scale cohort analysis, sST2 was determined as an independent biomarker for cardiovascular mortality in stable CAD patients. According to a study, long-term mortality has been estimated in stable CAD patients with a sST2 value of 24.6 ng/mL in blood plasma, and it has been reported that the risk of death increases two-fold in people above this value^[18].

According to the researches, only ST2 determined in plasma is insufficient for the correct diagnosis of the disease. The fact that the ST2 increasing mechanism that causes the formation of CAD is unknown and the absence of sST2 and gene expression studies in PF indicate that this study is important. In this study, it was aimed to determine ST2 and its gene expression level in PF of cardiovascular patients. The demographic characteristics of the patients included in the study for this purpose are presented in Table 7.

As a result of our study, compared to the control group, sST2 level of CAD patients was found to be 8.5 times higher in plasma and 5.5 times in PF.

The sST2 expression results were analyzed statistically both in the PF and plasma of the patient group and in the plasma of the

control group. The statistical results of the mean average (AVG) cycle threshold (Ct), AVG Delta Ct, $2^{\Delta(-AVG.(Delta(Ct))}$, fold change, and P-values obtained from sST2 are presented in Table 8. As a result of statistical analysis, significant changes were found in the replicate $2^{\Delta(-Delta Ct)}$ values of sST2 gene expression in PF and plasma ($P < 0.001$).

In our study, the increase in sST2 gene expression and sST2 level in CAD plasma and PF supports the prediction that sST2 value > 35 ng/ml increases the risk of CVD. This is actually an expected situation. In previous studies, it was noted that increased sST2 level, which inhibits the cardioprotective effect of IL33, is a biomarker in the diagnosis of CVD. High level of sST2 in the patient group may play an active role in the progression of the disease by contributing to the pathophysiologic mechanism in patients with CAD. While there are many tissues that increase the circulating level of sST2, the most important factor that increases the level of sST2 in PF is cardiovascular tissues, because sST2 release to PF can only be from the closest heart tissue.

In many previous studies, although sST2 release is known to increase in heart and lung diseases, its release focus and rate are still unknown^[19]. There are studies showing that sST2 protein is secreted after vascular endothelial cells IL-1b, tumor necrosis factor, and phorbol ester stimulation in response to an inflammatory stimulus^[20]. In some studies, it was found that there is an increase in sST2 in plasma of patients with pulmonary artery hypertension^[21]. In another study, it was reported that human peripheral blood cells and mast cells derived from mouse bone marrow and stimulated subsequently produce soluble sST2^[22].

Limitations

Our study has some limitations. This is a retrospective non-randomized study and it included a limited number of patients. Since PF cannot be obtained from healthy people, ST2 analysis could not be performed on them. Our main interest was to

Table 7. Demographic characteristics of the control and patient groups included in the study.

Variable	Control	Patients
Gender (female/male, %)	12/28 (30%/70%)	13/27 (32.5%/67.5%)
Age (years)	63	61

Table 8. Statistical results of gene expression of sST2 in PF and plasma.

sST2	Average Ct			Average Delta (Ct)			$2^{\Delta(-AVG.(Delta(Ct))}$			Fold change	P-value
	PF	CAD plasma	Control plasma	PF	CAD plasma	Control plasma	PF	CAD plasma	Control plasma		
	13.33±0.59	19.69±2.88	2.26±0.34	0.790741	0.39037	0.7450	0.57805	131.073	0.1506	22675	0.000265

AVG Delta (Ct)=(Ct (GOI) - AVG Ct (HKG)).

AVG=average; CAD=coronary artery disease; PF=pericardial fluid; Ct=cycle threshold; sST2=soluble growth STimulation expressed gene 2

investigate the prognostic value of ST2 in this study. Differently designed studies about the pathophysiological processes are needed to elucidate the underlying mechanisms of our findings.

CONCLUSION

In conclusion, determining the sST2 level only in plasma may not give very clear results for the reasons mentioned. Therefore, in order to make a real prediction, determining sST2 in both plasma and PF is important for knowing the oscillation focus. The study is consistent with previous studies showing that the main source of increased sST2 is not only the myocardium. Before our study, it was assumed that one of the sST2 sources in humans is probably the pericardium. In our study, the detection of high amounts of sST2 levels and gene expression in the PF of the patient group confirmed this prediction. In addition, the increase in both the level of sST2 gene expression and its amount in PF supports the prediction that it can be used as a prognostic biomarker for heart disease. Knowing the sST2 release foci can contribute to the development of more effective treatment methods, such as suppressing sST2 expression with various treatments or preventing sST2-IL33 binding.

ACKNOWLEDGMENTS

We would like to thank Harran University for the instrumental support.

No Financial support.

No conflict of interest.

Authors' roles & responsibilities

RD	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
MP	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
MI	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
IK	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
ET	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
MSA	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
ÖG	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published

REFERENCES

- Zhao D, Liu J, Wang M, Zhang X, Zhou M. Epidemiology of cardiovascular disease in China: current features and implications. *Nat Rev Cardiol*. 2019;16(4):203-12. doi:10.1038/s41569-018-0119-4.
- Spodick DH. Microphysiology of the pericardium: substrate for intrapericardial therapeutics. *Herz*. 2000;25(8):720-3. doi:10.1007/pl0001988.
- Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity*. 2005;23(5):479-90. doi:10.1016/j.immuni.2005.09.015.
- Sanada S, Hakuno D, Higgins LJ, Schreiter ER, McKenzie AN, Lee RT. IL-33 and ST2 comprise a critical biomechanically induced and cardioprotective signaling system. *J Clin Invest*. 2007;117(6):1538-49. doi:10.1172/JCI30634.
- Caselli C. Inflammation in cardiac disease: focus on interleukin-33/ST2 pathway. *Inflammation and Cell Signaling*. 2014;1:e149. doi:10.14800/ics.149.
- Dieplinger B, Mueller T. Soluble ST2 in heart failure. *Clin Chim Acta*. 2015;443:57-70. doi:10.1016/j.cca.2014.09.021.
- Wang TJ, Wollert KC, Larson MG, Coglianese E, McCabe EL, Cheng S, et al. Prognostic utility of novel biomarkers of cardiovascular stress: the Framingham heart study. *Circulation*. 2012;126(13):1596-604. doi:10.1161/CIRCULATIONAHA.112.129437.
- Weinberg EO, Shimpo M, De Keulenaer GW, MacGillivray C, Tominaga S, Solomon SD, et al. Expression and regulation of ST2, an interleukin-1 receptor family member, in cardiomyocytes and myocardial infarction. *Circulation*. 2002;106(23):2961-6. doi:10.1161/01.cir.0000038705.69871.d9.
- Truong QA, Januzzi JL, Szymonifka J, Thai WE, Wai B, Lavender Z, et al. Coronary sinus biomarker sampling compared to peripheral venous blood for predicting outcomes in patients with severe heart failure undergoing cardiac resynchronization therapy: the BIOCRT study. *Heart Rhythm*. 2014;11(12):2167-75. doi:10.1016/j.hrthm.2014.07.007.
- Ciccione MM, Cortese F, Gesualdo M, Riccardi R, Di Nunzio D, Moncelli M, et al. A novel cardiac bio-marker: ST2: a review. *Molecules*. 2013;18(12):15314-28. doi:10.3390/molecules181215314.
- McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the task force for the diagnosis and treatment of acute and chronic heart failure 2012 of the European society of cardiology. Developed in collaboration with the heart failure association (HFA) of the ESC. *Eur Heart J*. 2012;33(14):1787-847. Erratum in: *Eur Heart J*. 2013;34(2):158. doi:10.1093/eurheartj/ehs104.
- Moore SA, Januzzi JL Jr. Found in translation soluble ST2 and heart disease. *J Am Coll Cardiol*. 2010;55(3):251-3. doi:10.1016/j.jacc.2009.08.049.
- Eggers KM, Armstrong PW, Califf RM, Simoons ML, Venge P, Wallentin L, et al. ST2 and mortality in non-ST-segment elevation acute coronary syndrome. *Am Heart J*. 2010;159(5):788-94. doi:10.1016/j.ahj.2010.02.022.
- Dieplinger B, Egger M, Poelz W, Haltmayer M, Mueller T. Long-term stability of soluble ST2 in frozen plasma samples. *Clin Biochem*. 2010;43(13-14):1169-70. doi:10.1016/j.clinbiochem.2010.05.019.
- Coglianese EE, Larson MG, Vasan RS, Ho JE, Ghorbani A, McCabe EL, et al. Distribution and clinical correlates of the interleukin receptor family member soluble ST2 in the Framingham heart study. *Clin Chem*. 2012;58(12):1673-81. doi:10.1373/clinchem.2012.192153.
- Januzzi JL Jr. ST2 as a cardiovascular risk biomarker: from the bench to the bedside. *J Cardiovasc Transl Res*. 2013;6(4):493-500. doi:10.1007/s12265-013-9459-y.
- Ky B, French B, McCloskey K, Rame JE, McIntosh E, Shahi P, et al. High-sensitivity ST2 for prediction of adverse outcomes in

- chronic heart failure. *Circ Heart Fail.* 2011;4(2):180-7. doi:10.1161/CIRCHEARTFAILURE.110.958223.
18. Dieplinger B, Egger M, Haltmayer M, Kleber ME, Scharnagl H, Silbernagel G, et al. Increased soluble ST2 predicts long-term mortality in patients with stable coronary artery disease: results from the Ludwigshafen risk and cardiovascular health study. *Clin Chem.* 2014;60(3):530-40. doi:10.1373/clinchem.2013.209858.
19. Mildner M, Storka A, Lichtenauer M, Mlitz V, Ghannadan M, Hoetzenecker K, et al. Primary sources and immunological prerequisites for sST2 secretion in humans. *Cardiovasc Res.* 2010;87(4):769-77. doi:10.1093/cvr/cvq104.
20. Bartunek J, Delrue L, Van Durme F, Muller O, Casselman F, De Wiest B, et al. Nonmyocardial production of ST2 protein in human hypertrophy and failure is related to diastolic load. *J Am Coll Cardiol.* 2008;52(25):2166-74. doi:10.1016/j.jacc.2008.09.027.
21. Carlomagno G, Messalli G, Melillo RM, Stanziola AA, Visciano C, Mercurio V, et al. Serum soluble ST2 and interleukin-33 levels in patients with pulmonary arterial hypertension. *Int J Cardiol.* 2013;168(2):1545-7. doi:10.1016/j.ijcard.2012.12.031.
22. Bandara G, Beaven MA, Olivera A, Gilfillan AM, Metcalfe DD. Activated mast cells synthesize and release soluble ST2-a decoy receptor for IL-33. *Eur J Immunol.* 2015;45(11):3034-44. doi:10.1002/eji.201545501.



This is an open-access article distributed under the terms of the Creative Commons Attribution License.